

Class number formulas via 2-isogenies of elliptic curves

Cam McLeman
University of Michigan-Flint

Christopher Rasmussen
Wesleyan University

Abstract

A classical result of Dirichlet shows that certain elementary character sums compute class numbers of quadratic imaginary number fields. We obtain analogous relations between class numbers and a weighted character sum associated to a 2-isogeny of elliptic curves.

1 Introduction

We begin by recalling a famous result of Dirichlet which calculates the class number of a quadratic imaginary number field of prime discriminant via a finite sum. Throughout, we let $p > 3$ denote a prime number. Let $(\frac{a}{p})$ denote the usual Legendre symbol on $\mathbb{F}_p^\times$:

$$\left(\frac{a}{p}\right) = \begin{cases} +1 & a \in \mathbb{F}_p^{\times 2} \\ -1 & a \notin \mathbb{F}_p^{\times 2}. \end{cases}$$

The symbol is extended to all of $\mathbb{Z}$ via the reduction map $\mathbb{Z} \rightarrow \mathbb{F}_p$, and the definition $(\frac{a}{p}) = 0$ when $(a, p) > 1$. We let h_p denote the class number of $\mathbb{Q}(\sqrt{-p})$. For notational convenience, we set

$$h_p^* := \begin{cases} 0 & p \equiv 1 \pmod{4} \\ h_p & p \equiv 3 \pmod{4}. \end{cases}$$

The following consequence of Dirichlet's class number formula (e.g., [2, Ch. 6]) is well-known:

Theorem 1 (Dirichlet). *For any prime $p > 3$,*

$$-\frac{1}{p} \sum_{x=1}^{p-1} x \left(\frac{x}{p}\right) = h_p^*. \quad (1)$$

This article has been accepted for publication in the Bulletin of the London Mathematical Society. However, it does not reflect any changes or corrections between acceptance and publication; the two versions may be different.

2000 Mathematics Subject Classification: 11G05 (primary), 11R29, 11T24, 11G20 (secondary)

This research was supported in part by the Van Vleck Fund at Wesleyan University.

We consider the following point of view for Dirichlet's result. The $\mathbb{F}_p$ -rational morphism $\phi: \mathbb{G}_m \rightarrow \mathbb{G}_m$ defined by $\phi(x) = x^2$ partitions the points of $\mathbb{G}_m(\mathbb{F}_p) \cong \mathbb{F}_p^\times$ into two sets – those that are the image of an $\mathbb{F}_p$ -rational point of the domain (i.e., the quadratic residues), and those that are not (the non-residues). The character $\chi_\phi := \left(\frac{\cdot}{p}\right)$ is now precisely the natural identification of the cokernel of ϕ with $\{\pm 1\}$ which makes the following sequence exact:

$$\mathbb{G}_m(\mathbb{F}_p) \xrightarrow{\phi} \mathbb{G}_m(\mathbb{F}_p) \xrightarrow{\chi_\phi} \{\pm 1\} \longrightarrow 0 .$$

Note that it is the properties of ϕ , not the underlying algebraic group $\mathbb{G}_m$, which allow this construction. This paper demonstrates that an analogous procedure, arising from a different morphism of algebraic groups, yields new character sums with similar arithmetic properties. Let τ be a degree 2 isogeny of elliptic curves defined over $\mathbb{F}_p$. We define a weighted character sum S_τ , analogous to the sum appearing in (1). The quantity S_τ is shown to be divisible by p , and a strong relationship between S_τ and h_p^* is established.

Remark. Throughout this article, we study sums of the form $\sum g(x)\chi(x)$, where *a priori* the values of $g(x)$ lie in the finite field $\mathbb{F}_p$. We use the following convention, so as to view the value of the sum as an integer: Each summand is the scaling of the character value $\chi(x) \in \mu_2(\mathbb{C}) = \{\pm 1\}$ by the unique integral lift of $g(x)$ in the range $[0, p)$. For clarity, we use braces $\{\cdot\}$ to denote the lifting $\mathbb{F}_p \rightarrow \mathbb{Z} \cap [0, p)$ explicitly.

In some proofs, it will be convenient to view sums of the form $\sum_{x=0}^{p-1}$ interchangeably as sums over $\mathbb{F}_p$ or as sums over the range $[0, p)$ of integers. Consequently, there will occasionally be a mild abuse of notation – for example, given $a \in \mathbb{Z}$, we may write $\{a\}$ to mean $\{\bar{a}\}$, where $\bar{a}$ is the reduction of a mod p .

CM Example

We begin with two typical results, both special cases of the main theorem. Consider first the elliptic curve

$$E/\mathbb{Q}: \quad y^2 = (x+2)(x^2-2), \quad (2)$$

which possesses complex multiplication by $\mathbb{Z}[\sqrt{-2}]$. As this ring possesses elements of absolute norm 2, there exist endomorphisms of degree 2 on E . These endomorphisms admit reductions defined over $\mathbb{F}_p$ whenever p is a prime of good and ordinary reduction (equivalently, $\left(\frac{-2}{p}\right) = 1$). For a specific choice of τ (see §4), we set

$$\chi_\tau(P) = \begin{cases} +1 & P \in \tau(E(\mathbb{F}_p)) \\ -1 & P \notin \tau(E(\mathbb{F}_p)) \end{cases}. \quad (3)$$

This defines a character on $E(\mathbb{F}_p)$, and the following sequence is exact:

$$E(\mathbb{F}_p) \xrightarrow{\tau} E(\mathbb{F}_p) \xrightarrow{\chi_\tau} \mu_2 \longrightarrow 0 .$$

The following is a consequence of the main theorem:

Proposition 2. *Let $p > 3$ be a prime of good and ordinary reduction for the elliptic curve E given in (2). Then*

$$-\frac{1}{p} \sum_{\substack{P \in E(\mathbb{F}_p) \\ P \neq \infty}} \{x(P)\} \chi_\tau(P) = h_p^*. \quad (4)$$

Remark. It is not *a priori* clear that the sum in (4) (nor the sum in (1), for that matter) is divisible by p . We note that the two sums appearing in (1) and (4) are *not* the same expressions, even though they both compute h_p^* . This is immediate from the observation that E and $\mathbb{G}_m$ need not have the same number of points over $\mathbb{F}_p$. Here is an explicit example: When $p = 11$, one has

$$E(\mathbb{F}_{11}) = \{\infty, (7, \pm 4), (8, \pm 2), (9, 0)\}, \quad \tau(E(\mathbb{F}_{11})) = \{\infty, (7, \pm 4)\}.$$

Hence, (4) evaluates as

$$-\frac{1}{11}(7 + 7 - 8 - 8 - 9) = 1,$$

whereas the classical expression (1) yields

$$-\frac{1}{11}(1 - 2 + 3 + 4 + 5 - 6 - 7 - 8 + 9 - 10) = 1.$$

Non-CM Example

This connection between weighted character sums and class numbers is not unique to isogenies arising from complex multiplication. Consider the elliptic curves

$$\begin{aligned} E_1/\mathbb{Q}: \quad y^2 &= x^3 + 2x^2 - x, \\ E_2/\mathbb{Q}: \quad y^2 &= x^3 - 4x^2 + 8x, \end{aligned}$$

and the following isogeny of degree two:

$$\tau: E_1 \rightarrow E_2, \quad \tau(x, y) = \left(\frac{y^2}{x^2}, -\frac{y(1+x^2)}{x^2} \right).$$

The curves E_1 and E_2 have good reduction away from 2. For any odd prime p , τ induces an isogeny between the reductions of E_1 and E_2 over $\mathbb{F}_p$, and this morphism is in fact $\mathbb{F}_p$ -rational. Hence, there exists a homomorphism $\tau: E_1(\mathbb{F}_p) \rightarrow E_2(\mathbb{F}_p)$. As in the previous example, we consider the character

$$\chi_\tau: E_2(\mathbb{F}_p) \rightarrow \{\pm 1\},$$

where $\chi_\tau(P) = +1$ if and only if $P \in \tau(E_1(\mathbb{F}_p))$. Again we find a strong relationship between h_p^* and the weighted character sum

$$S_\tau := \sum_{\substack{P \in E_2(\mathbb{F}_p) \\ P \neq \infty}} \{x(P) - 2\} \chi_\tau(P).$$

The following is a special case of the main theorem (see §4, Example 1):

Proposition 3. *With E_1 , E_2 , and τ as above, and any prime $p > 3$, we have $-\frac{1}{p}S_\tau = h_p^*$.*

Main result

The main theorem generalizes the previous examples, which each relate h_p^* to a weighted character sum. Suppose $E_1/\mathbb{Q}$, $E_2/\mathbb{Q}$ are elliptic curves and $\tau: E_1 \rightarrow E_2$ is a $\mathbb{Q}$ -rational 2-isogeny. For any prime p of good reduction, there is an $\mathbb{F}_p$ -rational isogeny $\tau_p: E_1/\mathbb{F}_p \rightarrow E_2/\mathbb{F}_p$. Hence, τ induces a family of isogenies $\{\tau_p\}$, indexed by the primes of good reduction. To each of these isogenies, there is an associated character $\chi_\tau = \chi_{\tau,p}$, and an associated weighted character sum $S_{\tau,p}$ (defined below). As in the above examples, the quotient $-\frac{1}{p}S_{\tau,p}$ always approximates h_p^* well, in the sense that there is an absolute bound for the error as p varies among all primes of good reduction.

Concretely, let $a, b \in \mathbb{Z}$, and let $E_1/\mathbb{Q}$ and $E_2/\mathbb{Q}$ be the elliptic curves given by the following Weierstrass models:

$$\begin{aligned} E_1: \quad y^2 &= x^3 + ax^2 + bx, \\ E_2: \quad y^2 &= x^3 - 2ax^2 + (a^2 - 4b)x \end{aligned} \tag{5}$$

Let $\tau: E_1 \rightarrow E_2$ be the explicit isogeny given in (7).

Main Theorem. Let E_1 , E_2 , and τ be as above, and let $p > 3$ be a prime of good reduction for E_1 and E_2 .

(a) For all such p , the weighted character sum

$$S_{\tau,p} := \sum_{\substack{P \in E_2(\mathbb{F}_p) \\ P \neq \infty}} \{x(P) - a\} \chi_\tau(P)$$

is divisible by p .

(b) $S_{\tau,p}$ approximates $-ph_p^*$ in the following sense: the quantity

$$R_{a,b}(p) = -\frac{1}{p}S_{\tau,p} - h_p^*$$

is bounded in absolute value by a constant C_τ , independent of p .

(c) If there exists one $p > |a|$ such that $R_{a,b}(p) = 0$, then there exists a set of primes of positive density for which $S_{\tau,p} = h_p^*$, determined by explicit congruence conditions.

The remainder of the paper is organized as follows. In §2, we collect relevant facts about 2-isogenies over finite fields, and compute the cokernel character χ_τ in terms of the Tate pairing. In §3, we prove the main theorem. The proof combines classical techniques for evaluating character sums with formulas deduced from the Tate pairing. In §4, we extend the theorem to isogenies not specifically of the form (7), in particular the degree 2 endomorphisms existing on elliptic curves with complex multiplication by $\mathbb{Z}[\sqrt{-1}]$, $\mathbb{Z}[\sqrt{-2}]$, and $\mathbb{Z}[\sqrt{-7}]$. An analogous result for the dual isogeny $\hat{\tau}$ of (7) is given in the Appendix.

2 Preliminaries

Our strategy for the proof of the Main Theorem will be to convert the characters in the introduction into explicitly computable Legendre symbols. For this, we recall some facts about isogenies of degree 2 and the mechanics of the Tate pairing attached to an isogeny.

Degree 2 isogenies

Let K be an arbitrary field with $\text{char } K \neq 2$, and let E_1 and E_2 be elliptic curves defined over K . Let $\phi: E_1 \rightarrow E_2$ be an isogeny of degree 2 defined over K . Necessarily, this implies that both E_1 and E_2 possess K -rational points of order 2 (generating the kernels of ϕ and the dual $\hat{\phi}$, respectively). Hence, by appropriate changes of coordinates, E_1 and E_2 are isomorphic over K to elliptic curves E'_1 and E'_2 , respectively, which each possess Weierstrass equations of the form $y^2 = f_i(x)$, with $f_1(0) = f_2(0) = 0$. In fact, we may always simultaneously choose isomorphisms $\alpha_i: E_i \rightarrow E'_i$ such that:

- (i) E'_1 has the Weierstrass model $y^2 = x^3 + ax^2 + bx$ with $a, b \in K$;
- (ii) E'_2 has the Weierstrass model $y^2 = x^3 - 2ax^2 + rx$, with $r = a^2 - 4b$;
- (iii) There exists a K -rational 2-isogeny $\tau: E'_1 \rightarrow E'_2$ such that the following diagrams commute:

$$\begin{array}{ccc} E_1 & \xrightarrow{\alpha_1} & E'_1 \\ \phi \downarrow & & \downarrow \tau \\ E_2 & \xrightarrow{\alpha_2} & E'_2 \end{array} \quad \begin{array}{ccc} E_2 & \xrightarrow{\alpha_2} & E'_2 \\ \hat{\phi} \downarrow & & \downarrow \hat{\tau} \\ E_1 & \xrightarrow{\alpha_1} & E'_1 \end{array} \quad (6)$$

The explicit formulas for τ and its dual are well-known [7, III.4.5]:

$$\tau(x, y) = \left(\frac{y^2}{x^2}, \frac{y(b - x^2)}{x^2} \right), \quad \hat{\tau}(x, y) = \left(\frac{y^2}{4x^2}, \frac{y(r - x^2)}{8x^2} \right). \quad (7)$$

Placing the isogeny in this form will simplify the computations involving the Tate pairing. We will show how to treat more general 2-isogenies in Section 4.

The Tate pairing

Let ℓ and p be prime numbers such that $p \equiv 1 \pmod{\ell}$, and let E_1 and E_2 be elliptic curves defined over $\mathbb{F}_p$. Let $\tau: E_1 \rightarrow E_2$ be an isogeny of degree ℓ defined over $\mathbb{F}_p$, and let $\hat{\tau}$ denote the dual isogeny. Let us assume that $E_1[\tau] \subseteq E_1(\mathbb{F}_p)$ and $E_2[\hat{\tau}] \subseteq E_2(\mathbb{F}_p)$, and let T_1 and T_2 generate the groups $E_1[\tau]$ and $E_2[\hat{\tau}]$, respectively. The Tate pairing associated to τ is a function

$$\psi_\tau: \frac{E_2(\mathbb{F}_p)}{\tau(E_1(\mathbb{F}_p))} \times E_2[\hat{\tau}] \longrightarrow \mathbb{F}_p^\times / \mathbb{F}_p^{\times \ell}$$

which is bilinear and non-degenerate on the left. (See, for example, [7, X.1.1], which develops the properties of $\psi_{[m]}$. The proofs for ψ_τ are essentially identical.) We can explicitly compute ψ_τ as follows. The points T_1 and T_2 are of exact order ℓ , as is any $R \in \tau^{-1}(T_2)$. Thus, there exist functions f on E_2 and g on E_1 whose divisors are:

$$\begin{aligned} (f) &= \ell(T_2) - \ell(\infty), \\ (g) &= \tau^*((T_2) - (\infty)) = \sum_{i=0}^{\ell-1} (R + iT_1) - \sum_{i=0}^{\ell-1} (iT_1). \end{aligned} \quad (8)$$

Scaling either f or g by a constant if necessary, we have an equality of functions $f \circ \tau = g^\ell$. For a point $S \in E_2(\mathbb{F}_p)$, let $[S]$ denote its image in $E_2(\mathbb{F}_p)/\tau(E_1(\mathbb{F}_p))$. If $S \in E_2(\mathbb{F}_p)$ and $[S] \neq [T_2], [\infty]$, we define

$$\psi_\tau([S], T_2) := f(S) \pmod{\mathbb{F}_p^{\times \ell}}.$$

In case $[S] \in \{[T_2], [\infty]\}$, we choose a point $Q \notin \{T_2, \infty\}$ on E_2 and set

$$\psi_\tau([S], T_2) := \frac{f(S+Q)}{f(Q)} \pmod{\mathbb{F}_p^{\times \ell}}.$$

As $E_2[\hat{\tau}]$ is generated by T_2 , we may recover the entire pairing by bilinearity, since we have $\psi_\tau([S], kT_2) = \psi_\tau([S], T_2)^k$. Thus, the following definition is complete:

$$\psi_\tau([S], kT_2) := \begin{cases} f(S)^k & [S] \notin \{[T_2], [\infty]\} \\ \left(\frac{f(S+Q)}{f(Q)}\right)^k & [S] \in \{[T_2], [\infty]\}. \end{cases}$$

Formulas for 2-isogenies

We now specialize to the case of the degree 2 isogenies τ and $\hat{\tau}$ given in (7).

Definition 4. Define a character $\chi_\tau: E_2(\mathbb{F}_p) \rightarrow \mu_2$ by

$$\chi_\tau(P) = \begin{cases} +1 & P \in \tau(E_1(\mathbb{F}_p)) \\ -1 & P \notin \tau(E_1(\mathbb{F}_p)). \end{cases}$$

We compute this character explicitly in terms of the pairing ψ_τ . This allows us to replace χ_τ with expressions involving the Legendre symbol, and so evaluate the weighted character sum S_τ from the introduction. We let T_1 and T_2 respectively denote the point $(0,0) \in E_1(\mathbb{F}_p)$, which generates $E_1[\tau]$, and the point $(0,0) \in E_2(\mathbb{F}_p)$, which generates $E_2[\hat{\tau}]$.

Proposition 5. Suppose $\tau: E_1 \rightarrow E_2$ is of the form (7), and let $P = (x, y) \in E_2(\mathbb{F}_p)$. Then

$$\chi_\tau(P) = \psi_\tau([P], T_2) = \begin{cases} \left(\frac{x}{p}\right) & [P] \neq [T_2], [\infty] \\ \left(\frac{y}{p}\right) & [P] = [T_2] \\ 1 & [P] = [\infty], \end{cases}$$

where in the second equality we canonically identify $\mathbb{F}_p^\times/\mathbb{F}_p^{\times 2}$ with μ_2 via the Legendre symbol.

Proof. For the statement that $\chi(\cdot) = \psi_\tau(\cdot, T_2)$, we must show that a point $P \in E_2(\mathbb{F}_p)$ is in the image of τ if and only if $\psi_\tau([P], T_2) = 1$. By bilinearity, $\psi_\tau([P], kT_2) = \psi_\tau([P], T_2)^k$. As $E_2[\hat{\tau}]$ is generated by T_2 , P pairs trivially with T_2 if and only if it pairs trivially with every element of $E_2[\hat{\tau}]$. By the left non-degeneracy of ψ_τ , this occurs if and only if $[P]$ represents the trivial class of $E_2(\mathbb{F}_p)/\tau(E_1(\mathbb{F}_p))$, i.e., P is in the image of τ .

To prove the second equality, we now compute $\psi_\tau([P], T_2)$ explicitly. Functions f and g whose divisors are given in (8) are $f(x, y) = x$, $g(x, y) = \frac{y}{x}$. No scaling is necessary, as we have

$$f(\tau(P)) = x(\tau(P)) = \frac{y^2}{x^2} = g^2(P).$$

Thus, $\psi_\tau([P], T_2)$ is a square in $\mathbb{F}_p^\times$ if and only if $(\frac{x}{p}) = 1$. This proves the second equality when $[P] \neq [T_2], [\infty]$. The result is trivial for $[P] = [\infty]$. It remains to prove $\psi_\tau([T_2], T_2) = (\frac{r}{p})$. Notice that $T_2 \in \tau(E_1(\mathbb{F}_p))$ if and only if $E_1[2] \subseteq E_1(\mathbb{F}_p)$. From this we see

$$\begin{aligned} \psi_\tau([T_2], T_2) = +1 &\iff E_1[2] \subseteq E_1(\mathbb{F}_p) \\ &\iff x^2 + ax + b \text{ splits in } \mathbb{F}_p[x] \\ &\iff r = a^2 - 4b \equiv \square \pmod{p} \\ &\iff \left(\frac{r}{p}\right) = +1. \end{aligned}$$

□

Remark. In fact, $[T_2] = [\infty]$ if and only if $(\frac{r}{p}) = 1$.

Dual Isogeny Formulas

Finally, we note that we can provide an equally explicit result for the pairing attached to the dual $\hat{\tau}$. The cokernel character $\chi_{\hat{\tau}}$ is defined by

$$\chi_{\hat{\tau}}(P) = \begin{cases} +1 & P \in \hat{\tau}(E_2(\mathbb{F}_p)) \\ -1 & P \notin \hat{\tau}(E_2(\mathbb{F}_p)). \end{cases}$$

Let $\hat{f}$ and $\hat{g}$ be the functions on E_1 whose divisors are

$$\begin{aligned} (\hat{f}) &= 2(T_1) - 2(\infty), \\ (\hat{g}) &= \hat{\tau}^*((T_1) - (\infty)). \end{aligned}$$

For example, take $\hat{f}(x, y) = x$, and $\hat{g}(x, y) = \frac{y}{2x}$. Then the associated pairing

$$\psi_{\hat{\tau}}: \frac{E_1(\mathbb{F}_p)}{\hat{\tau}(E_2(\mathbb{F}_p))} \times E_1[\tau] \longrightarrow \frac{\mathbb{F}_p^\times}{\mathbb{F}_p^{\times 2}}$$

may be computed via

$$\psi_{\hat{\tau}}([S], kT_1) = \begin{cases} \hat{f}(S)^k & [S] \notin \{[T_1], [\infty]\} \\ \left(\frac{\hat{f}(S+Q)}{\hat{f}(Q)}\right)^k & [S] \in \{[T_1], [\infty]\}. \end{cases}$$

Proposition 6. *For any $P = (x, y) \in E_1(\mathbb{F}_p)$,*

$$\chi_{\hat{\tau}} = \psi_{\hat{\tau}}([P], T_1) = \begin{cases} \left(\frac{x}{p}\right) & [P] \neq [T_1], [\infty] \\ \left(\frac{y}{p}\right) & [P] = [T_1] \\ 1 & [P] = [\infty]. \end{cases}$$

Proof. The argument parallels Proposition 5 exactly. $\square$

3 Weighted character sums

There are well-established connections between arithmetic data and character sums arising from elliptic curves. If $y^2 = f(x)$ is an integral model for an elliptic curve over $\mathbb{Q}$ with complex multiplication, and p is a prime of good reduction, then the work of Deuring [3] demonstrates that character sums of the form

$$\sum_{x=1}^{p-1} \left(\frac{f(x)}{p}\right)$$

can be computed in terms of the trace of Frobenius and the splitting of p in the endomorphism ring of the curve. Similar results have been established by many different authors. We mention, for example, the works of Williams [10], Joux-Morain [4], and Padma-Venkataraman [6], which each take slightly different approaches to such character sums, consolidate many previous results, and contain comprehensive bibliographies.

We briefly contrast these character sums to those contained in the present paper. First, the character χ_{τ} is determined by an isogeny, not an elliptic curve. Second, the terms of the sum are weighted by a non-trivial integer-valued function. To the authors' knowledge, these sums have not been extensively studied.

We recall our convention to use $\{\cdot\}$ to denote the lifting $\mathbb{F}_p \rightarrow \mathbb{Z} \cap [0, p)$, and introduce a second convention that a primed sum over the points on an elliptic curve will exclude the point at infinity on that curve. Under these conventions, we seek to evaluate the sum

$$\sum'_{P \in E_2(\mathbb{F}_p)} \{x(P) - a\} \chi_{\tau}(P), \tag{9}$$

where τ is the isogeny given in (7). Here, we are taking E_1 , E_2 , and τ to be defined over $\mathbb{F}_p$.

This is necessary if we hope to obtain interesting results, since trivially we have $\sum_P \chi_{\tau}(P) = 0$.

We first evaluate a useful character sum. For a given prime p and $k \in \mathbb{Z}$ such that $p \nmid k$, we define

$$\delta_k := \frac{1}{2} \left(1 + \left(\frac{k}{p} \right) \right) = \begin{cases} 1 & k \equiv \square \pmod{p}, \\ 0 & k \not\equiv \square \pmod{p}. \end{cases}$$

Lemma 7. *For any integer k relatively prime to p ,*

$$\sum_{u=1}^{p-1} u \left(\frac{u^2 + k}{p} \right) = -p\delta_k. \quad (10)$$

Proof. Let S be the sum. Then by substituting $u \mapsto p - u$, we find

$$S = \sum_{u=1}^{p-1} (p - u) \left(\frac{(p - u)^2 + k}{p} \right) = p \sum_{u=1}^{p-1} \left(\frac{u^2 + k}{p} \right) - S.$$

Therefore,

$$\begin{aligned} \frac{2S}{p} &= \sum_{u=1}^{p-1} \left(\frac{u^2 + k}{p} \right) = - \left(\frac{k}{p} \right) + \sum_{u=0}^{p-1} \left(\frac{u^2 + k}{p} \right) \\ &= - \left(\frac{k}{p} \right) - p + \sum_{u=0}^{p-1} \left[1 + \left(\frac{u^2 + k}{p} \right) \right]. \end{aligned} \quad (11)$$

Let $C/\mathbb{F}_p$ be the conic $u^2 + kw^2 = v^2$. The final term in (11) counts the number of $\mathbb{F}_p$ -rational points on C within the affine region $w \neq 0$. Since $(p, k) = 1$, the conic is birational to $\mathbb{P}^1$ and has $p + 1$ points. Exactly two of these, $(1 : \pm 1 : 0)$, lie on $w = 0$, so the affine region contains $p - 1$ points. Thus

$$S = \frac{p}{2} \left[- \left(\frac{k}{p} \right) - p + (p - 1) \right] = -\frac{p}{2} \left(1 + \left(\frac{k}{p} \right) \right) = -p\delta_k,$$

which completes the proof. $\square$

Remark. Sums of the form (10) will appear in the proof below, with $u = x - a$. This motivates the choice $g = x(P) - a$ as the weight in (9). We mention in passing that this function g has an interesting geometric description: If $Q_i = (\alpha_i, 0)$ are the 2-torsion points on E_2 which are not in the kernel of $\hat{\tau}$, then $a = \frac{1}{2}(\alpha_1 + \alpha_2)$. That is, we may think of $g = 0$ as the unique vertical line which intersects the x -axis at a point equidistant to both Q_1 and Q_2 .

As always, let $p > 3$ be prime. We have already seen at the start of §2 that we may transform any $\mathbb{F}_p$ -rational 2-isogeny into the isogeny τ of (7). Then the isogeny τ is between the curves

$$\begin{aligned} E_1: \quad & y^2 = f_1(x) = x^3 + ax^2 + bx, & a, b \in \mathbb{F}_p, \\ E_2: \quad & y^2 = f_2(x) = x^3 - 2ax^2 + rx, & r = a^2 - 4b. \end{aligned}$$

Recall that $T := (0, 0) \in E_2(\mathbb{F}_p)$ generates the kernel of $\hat{\tau}$. Our goal is to evaluate the sum

$$S_\tau := \sum'_{P \in E_2(\mathbb{F}_p)} \{x(P) - a\} \chi_\tau(P). \quad (12)$$

For convenience, we define an error term:

$$R_{a,b} := \delta_{-b} - \sum_{x=1}^{\{a\}-1} \left(\frac{x}{p}\right).$$

Proposition 8. *Let $p > 3$ be prime, and E_1 , E_2 , τ , and S_τ as above. Then S_τ is divisible by p and*

$$-\frac{1}{p}S_\tau = h_p^* + R_{a,b}. \quad (13)$$

Further, $|R_{a,b}| \leq \{a\}$.

Proof. The bound on $R_{a,b}$ is immediate. As h_p^* and $R_{a,b}$ are integers, it remains only to establish (13). Applying Proposition 5, we have:

$$\begin{aligned} S_\tau &= \sum'_{P \in E_2(\mathbb{F}_p)} (x(P) - a) \chi_\tau(P) \\ &= \{x(T) - a\} \chi_\tau(T) + \sum_{\substack{P \in E_2(\mathbb{F}_p) \\ P \neq T, \infty}} \{x(P) - a\} \left(\frac{x(P)}{p}\right) \\ &= \{p - a\} \left(\frac{r}{p}\right) + \sum_{x=1}^{p-1} \{x - a\} \left(\frac{x}{p}\right) \left(1 + \left(\frac{x^3 - 2ax^2 + rx}{p}\right)\right) \\ &= \{p - a\} \left(\frac{r}{p}\right) + \sum_{x=1}^{p-1} \{x - a\} \left(\frac{x}{p}\right) + \sum_{x=1}^{p-1} \{x - a\} \left(\frac{(x-a)^2 - 4b}{p}\right) \\ &= \sum_{x=1}^{p-1} \{x - a\} \left(\frac{x}{p}\right) + \sum_{x=0}^{p-1} \{x - a\} \left(\frac{(x-a)^2 - 4b}{p}\right). \end{aligned}$$

We evaluate these two sums in turn. First,

$$\begin{aligned} \sum_{x=1}^{p-1} \{x - a\} \left(\frac{x}{p}\right) &= \sum_{x=1}^{\{a\}-1} (p + \{x\} - \{a\}) \left(\frac{x}{p}\right) + \sum_{x=\{a\}}^{p-1} (\{x\} - \{a\}) \left(\frac{x}{p}\right) \\ &= p \sum_{x=1}^{\{a\}-1} \left(\frac{x}{p}\right) + \sum_{x=1}^{p-1} x \left(\frac{x}{p}\right) - a \sum_{x=1}^{p-1} \left(\frac{x}{p}\right) \\ &= p \sum_{x=1}^{\{a\}-1} \left(\frac{x}{p}\right) - ph_p^*. \end{aligned}$$

We note that the second sum is unchanged by making the substitution $u = x - a$. Thus,

$$\sum_{x=0}^{p-1} \{x - a\} \left(\frac{(x-a)^2 - 4b}{p} \right) = \sum_{u=0}^{p-1} \{u\} \left(\frac{u^2 - 4b}{p} \right) = -p\delta_{-b},$$

by Lemma 7. Combining the two sums, we have

$$S_\tau = -ph_p^* - p\delta_{-b} + p \sum_{x=1}^{\{a\}-1} \left(\frac{x}{p} \right),$$

which gives (13). $\square$

The Main Theorem can be viewed as the global version of Proposition 8. Consider elliptic curves $E_1/\mathbb{Q}$ and $E_2/\mathbb{Q}$ with respective Weierstrass models

$$\begin{aligned} E_1: y^2 &= x^3 + ax^2 + bx, \\ E_2: y^2 &= x^3 - 2ax^2 + (a^2 - 4b)x, \end{aligned}$$

with $a, b \in \mathbb{Z}$. As always, we let $\tau: E_1 \rightarrow E_2$ be the explicit $\mathbb{Q}$ -rational 2-isogeny given in (7).

Theorem 9. *Let $\tau: E_1 \rightarrow E_2$ be as above, and let $p > 3$ be a prime of good reduction.*

(a) *For all such p , the weighted character sum*

$$S_{\tau,p} := \sum'_{P \in E_2(\mathbb{F}_p)} \{x(P) - a\} \chi_\tau(P)$$

is divisible by p .

(b) *$S_{\tau,p}$ approximates $-ph_p^*$ in the following sense: the quantity*

$$R_{a,b}(p) = -\frac{1}{p}S_{\tau,p} - h_p^*$$

is bounded in absolute value by a constant C_τ , independent of p .

(c) *If there exists one $p > |a|$ such that $R_{a,b}(p) = 0$, then there exists a set of primes of positive density for which $S_{\tau,p} = h_p^*$, determined by explicit congruence conditions.*

Proof. Part (a) is precisely Proposition 8 applied to any prime of good reduction. For part (b), it is enough to bound the number of terms in $R_{a,b}$ independently of p . If $a > 0$, this is obvious; the sum has exactly $\{a\}$ terms, and $\{a\} \leq a$. When $a = 0$ we trivially have $C_\tau = 1$. Now suppose $a < 0$. It is no obstruction to assume $p > |a|$, and so $\{a\} = p + a$. We have

$$R_{a,b} = \delta_{-b} - \sum_{x=1}^{\{a\}-1} \left(\frac{x}{p} \right) = \delta_{-b} - \sum_{x=1}^p \left(\frac{x}{p} \right) + \sum_{x=\{a\}}^p \left(\frac{x}{p} \right) = \delta_{-b} + \sum_{x=p+a}^p \left(\frac{x}{p} \right),$$

and so $|R_{a,b}| \leq |a| + 2$.

For part (c), suppose $p_0 > |a|$ is a prime for which $R_{a,b}(p_0) = 0$. So $p = p_0$ is a solution to:

$$\delta_{-b} = \sum_{x=1}^{\{a\}-1} \left(\frac{x}{p}\right). \quad (14)$$

Now, among $p > |a|$, the individual terms of the sum (14) never vanish, and so the sum is periodic as a function of p , with respect to some sufficiently large modulus, e.g., $N = 4G$, where G is the least common multiple of $\{2, 3, \dots, |a| - 1\}$. Thus, every prime in the sequence $\{p_0 + kN\}$ also satisfies $R_{a,b}(p) = 0$. $\square$

The analogous result for the dual isogeny $\hat{\tau}$ is proved in the appendix.

Remark. It is natural to ask what happens when the hypothesis in part (c) does not hold. It is possible that the error term is zero for only finitely many p . For example, when $(a, b) = (9, -1)$, the error term vanishes only for $p = 7$. It is even possible that the error term is never zero (e.g., Example 2 below). Indeed, note that by the periodicity modulo N , the error term is never zero if it is non-zero for all $p < N$.

Remark. For fixed a and varying p , it is not hard to argue that the uniform bounds constructed in the proof of Theorem 9 are the best possible. However, for a *particular* value of p , better bounds certainly exist. For example, if $a = O(\log p)$, then the estimates of Pólya-Vinogradov (and later improvements by Burgess) may offer substantial improvement. For details, see for example [5, §9.4].

Examples

We consider a few examples for illustration. In all cases, the selection of values (a, b) determines elliptic curves E_1, E_2 by (5) and an isogeny τ by (7).

Example 1. Set $(a, b) = (2, -1)$. Then we have $R_{a,b}(p) = 0$ for all primes $p > 3$. Hence, $-\frac{S_{\tau}}{p} = h_p^*$ for all such p .

Example 2. Set $(a, b) = (3, -1)$. We find $R_{a,b} = -\left(\frac{2}{p}\right)$, which is non-zero for all $p > 3$.

Example 3. Set $(a, b) = (7, 2)$. Then, for $p \neq 5$ ($R_{a,b}(5) \neq 0$ by a separate calculation),

$$R_{a,b} = \delta_{-2} - \sum_{x=1}^6 \left(\frac{x}{p}\right) = 2 + \delta_{-2} + \left(\frac{2}{p}\right) + \left(\frac{3}{p}\right) + \left(\frac{5}{p}\right) + \left(\frac{6}{p}\right).$$

We wish to decide when this sum vanishes. By a parity argument, we must have $\delta_{-2} = 0$, and of the four remaining Legendre symbols, three must evaluate to -1 , and one to $+1$. Further, we cannot have $\left(\frac{5}{p}\right) = 1$. Otherwise, $R_{a,b} = 0$ implies $\left(\frac{2}{p}\right) = \left(\frac{3}{p}\right) = \left(\frac{6}{p}\right) = -1$, contradicting the multiplicativity of the Legendre symbol. The three choices for which of the other Legendre symbols is to be -1

each lead to congruence conditions easily determined via quadratic reciprocity. Namely, we find:

$$-\frac{S_\tau}{p} = h_p^* \iff p \equiv 17, 43, 67, 83, 107, \text{ or } 113 \pmod{120}.$$

4 Complex Multiplication

Having computed the character sums associated to 2-isogenies given in the specific form (7), we turn to the analogous computation for other 2-isogenies. The principal technical lemma is that the character sums are unaffected by applying a change of coordinates $x \mapsto x - \varepsilon$ to the curve. This will allow us to change the codomain of an arbitrary 2-isogeny to one of the form in (7), and allow us to compute the corresponding character sums. As a primary application, the elliptic curves with complex multiplication by $\sqrt{-1}$, $\sqrt{-2}$, or $\sqrt{-7}$ possess an endomorphism of degree 2, and in this section we will compute the associated weighted character sum. In particular, we deduce Proposition 2.

Suppose that $\phi: E'_1 \rightarrow E'_2$ is a degree 2-isogeny defined over $\mathbb{F}_p$ with $p > 3$. Then there exist $\mathbb{F}_p$ -isomorphisms $\alpha_i: E'_i \rightarrow E_i$ and an isogeny τ of the form (7) such that the left square of the following diagram commutes:

$$\begin{array}{ccccc} E'_1 & \xrightarrow{\phi} & E'_2 & \xrightarrow{\chi_\phi} & \mu_2 \\ \alpha_1 \downarrow & & \downarrow \alpha_2 & & \parallel \text{id} \\ E_1 & \xrightarrow{\tau} & E_2 & \xrightarrow{\chi_\tau} & \mu_2 \end{array} \quad (15)$$

Here, χ_ϕ and χ_τ are the characters attached ϕ and τ , respectively.

Lemma 10. *The right square of the above diagram commutes, i.e., $\chi_\phi(P') = \chi_\tau(\alpha_2(P'))$ for any $P' \in E'_2(\mathbb{F}_p)$.*

Proof. This is a simple diagram chase. If $\chi_\phi(P') = 1$, then there exists a point $Q' \in E'_1(\mathbb{F}_p)$ such that $\phi(Q') = P'$. Hence, $\alpha_2(P') = \tau(\alpha_1(Q'))$, and so $\chi_\tau(\alpha_2(P')) = 1$ also. This argument is easily reversed, and the result follows. $\square$

As a consequence, we demonstrate the weighted character sum

$$S_\phi := \sum'_{P \in E'_2(\mathbb{F}_p)} \{x(P) - \xi\} \chi_\phi(P)$$

equals the sum S_τ of (12), provided α_2 is of a particular form and ξ is chosen appropriately. Let E'_2 be the elliptic curve

$$y^2 = (x - \varepsilon)(x^2 - 2\mu x + \nu), \quad \varepsilon, \mu, \nu \in \mathbb{F}_p. \quad (16)$$

defined over $\mathbb{F}_p$. Let ϕ be an isogeny such that the kernel of the dual isogeny $\hat{\phi}$ is generated by $(\varepsilon, 0) \in E'_2(\mathbb{F}_p)$. Set $(a, b) = (\mu - \varepsilon, \frac{1}{4}(\mu^2 - \nu))$, and let E_1, E_2, τ be as in (7). Then the map

$$\alpha_2(x, y) := (x - \varepsilon, y) \quad (17)$$

has the property that $\hat{\tau} \circ \alpha_2$ and $\hat{\phi}$ have the same kernel. This guarantees the existence of a unique α_1 satisfying $\hat{\tau} \circ \alpha_2 = \alpha_1 \circ \hat{\phi}$, and in fact, α_1 completes the diagram (15). Now set $\xi = \varepsilon + a$, so that

$$S_\phi = \sum'_{P \in E'_2(\mathbb{F}_p)} \{x(P) - \varepsilon - a\} \chi_\phi(P).$$

Lemma 11. $S_\phi = S_\tau$.

Proof. Let p be a prime of good reduction. Applying the previous lemma, we have

$$\begin{aligned} S_\phi &= \sum'_{P \in E'_2(\mathbb{F}_p)} \{x(P) - \varepsilon - a\} \chi_\phi(P) \\ &= \sum'_{P \in E_2(\mathbb{F}_p)} \{x(\alpha_2^{-1}(P)) - \varepsilon - a\} \chi_\phi(\alpha_2^{-1}(P)) \\ &= \sum'_{P \in E_2(\mathbb{F}_p)} \{x(P) - a\} \chi_\tau(P) = S_\tau. \end{aligned} \quad (18)$$

□

We now address CM endomorphisms. For each endomorphism ϕ , we select isomorphisms α_i such that $\alpha_2 \circ \phi = \tau \circ \alpha_1$, as in diagram (15), and such that E'_2 and α_2 have the forms (16) and (17), respectively. Evaluation of S_ϕ now follows from the previous lemma.

4.1 CM by -1

Consider first the elliptic curve $E/\mathbb{Q}$: $y^2 = x^3 + x$, which has complex multiplication by $\mathbb{Z}[\sqrt{-1}]$, and possesses the degree 2 endomorphism

$$\phi: E \longrightarrow E, \quad \phi(x, y) = \left(\frac{x^2 + 1}{2ix}, \frac{y(x^2 - 1)}{(2i - 2)x^2} \right). \quad (19)$$

Let $E'/\mathbb{Q}$ be the elliptic curve with Weierstrass equation $y^2 = x^3 - \frac{1}{4}x$, and let $\tau: E' \rightarrow E$ be the isogeny corresponding to $(a, b) = (0, -\frac{1}{4})$. If α_2 is the identity and α_1 is the isomorphism

$$\alpha_1(x, y) = (u^{-2}x, u^{-3}y), \quad u = -(1 + i),$$

then $\alpha_2 \circ \phi = \tau \circ \alpha_1$. Suppose $p > 3$ is a prime of good reduction which splits in $\mathbb{Q}(i)$. Each of the morphisms in the above diagram has a well-defined reduction

mod p , and the diagram still commutes after reduction. Applying Lemma 10 and Theorem 9, we see

$$S_\phi = S_\tau = -ph_p^* - pR_{0, -\frac{1}{4}} = -p.$$

This proves:

Corollary 12. *Let $p > 3$ be a prime, and suppose $p \equiv 1 \pmod{4}$. Then*

$$-\frac{1}{p} \sum'_{P \in E(\mathbb{F}_p)} \{x(P)\} \chi_\phi(P) = 1.$$

4.2 CM by -2

Next, we compute the character sum associated to a degree 2 endomorphism on an elliptic curve with complex multiplication by $\mathbb{Z}[\sqrt{-2}]$. One such elliptic curve is

$$E/\mathbb{Q}: y^2 = (x+2)(x^2-2).$$

One degree 2 endomorphism on E is:

$$\phi(x, y) = \left(\frac{(x+2)^2 + 2}{-2(x+2)}, \frac{y((x+2)^2 - 2)}{2\sqrt{-2}(x+2)^2} \right). \quad (20)$$

(This may be derived from the formula given in [8, pg. 111], but note that we are using a different Weierstrass equation.) Let $p > 3$ be a prime that splits in $\mathbb{Q}(\sqrt{-2})$, so that $\left(\frac{-2}{p}\right) = 1$. Equivalently, p is congruent to 1 or 3 (mod 8). For such primes, there is a well-defined reduction for ϕ over $\mathbb{F}_p$, which we will also denote ϕ . We now prove Proposition 2. Set

$$S_\phi := \sum'_{P \in E(\mathbb{F}_p)} \{x(P)\} \chi_\phi(P).$$

Corollary 13. *For any prime $p > 3$ such that $p \equiv 1$ or $3 \pmod{8}$, $-\frac{1}{p}S_\phi = h_p^*$.*

Proof. Let $E_1/\mathbb{Q}$ and $E_2/\mathbb{Q}$ denote the elliptic curves corresponding to the choice $(a, b) = (2, \frac{1}{2})$ in (7); let τ denote the corresponding 2-isogeny. It is straightforward to find isomorphisms α_i with the requisite properties such that $\tau \circ \alpha_1 = \alpha_2 \circ \phi$. Hence, $S_\phi = S_\tau$. By Proposition 8, $-\frac{1}{p}S_\tau = h_p^* + R_{2, \frac{1}{2}} = h_p^*$, as $R_{2, \frac{1}{2}} = 0$ if $p \equiv 1, 3 \pmod{8}$. $\square$

4.3 CM by -7

Finally, we compute the character sum associated to a degree 2 endomorphism on a curve with complex multiplication by $\mathbb{Z}[\sqrt{-7}]$. We take the elliptic curve

$$E/\mathbb{Q}: y^2 = (x+7)(x^2-7x+14).$$

Let $p > 3$ be a prime which splits in $\mathbb{Q}(\sqrt{-7})$, so $\left(\frac{-7}{p}\right) = 1$. Set $\beta = \frac{1+\sqrt{-7}}{2}$. A degree 2 endomorphism on E is given explicitly by ([8, pg. 111]):

$$\phi(x, y) = \left(\beta^{-2} \left(x - \frac{7(1-\beta^4)}{x+\beta^2-2} \right), \beta^{-3} y \left(1 + \frac{7(1-\beta)^4}{(x+\beta^2-2)^2} \right) \right). \quad (21)$$

Let E_1 and E_2 be the elliptic curves

$$\begin{aligned} E_1/\mathbb{Q}(\sqrt{-7}): y^2 &= x^3 + ax^2 + bx, \\ E_2/\mathbb{Q}(\sqrt{-7}): y^2 &= x^3 - 2ax^2 + rx, \end{aligned}$$

where

$$a = \frac{3}{2}(\beta - 4), \quad b = \frac{7}{16}(3\beta + 14), \quad r = a^2 - 4b.$$

The curves E , E_1 , and E_2 are all isomorphic over $\mathbb{Q}(\sqrt{-7})$. Let α_1 and α_2 be the isomorphisms

$$\begin{aligned} \alpha_1: E &\rightarrow E_1, & (x, y) &\mapsto ((\beta - 1)^2 x + \beta + 3, (\beta - 1)^3 y), \\ \alpha_2: E &\rightarrow E_2, & (x, y) &\mapsto (x + 4 - \beta, y). \end{aligned}$$

As before, we have $\tau \circ \alpha_1 = \alpha_2 \circ \phi$, and each morphism reduces to an $\mathbb{F}_p$ -rational morphism on the reduced curves, as p splits in $\mathbb{Q}(\sqrt{-7})$. If we define

$$S_\phi := \sum'_{P \in E(\mathbb{F}_p)} \{x(P)\} \chi_\phi(P),$$

then $S_\phi = S_\tau$, and we find $-\frac{1}{p}S_\phi = h_p^* + R_{a,b}$.

Unlike all previous examples, however, we do not have a uniform bound for the error. Our bound is always in terms of $\{a\}$, but in this situation $a \notin \mathbb{Z}$. Hence, as a function of p , $\{a\}$ is not eventually constant, and the previous results bounding $R_{a,b}$ do not apply.

Appendix: A proof of a dual isogeny calculation

Let τ and $\hat{\tau}$ be the isogenies defined in (7), with $a, b \in \mathbb{F}_p$ for $p > 3$. Recall that we set $r = a^2 - 4b$. Let $\eta = \{\frac{a}{2}\}$, and define

$$\hat{R}_{a,b} := \delta_{-r} + \sum_{x=1}^{\eta} \left(\frac{-x}{p} \right).$$

We wish to evaluate

$$S_{\hat{\tau}} := \sum'_{P \in E_1(\mathbb{F}_p)} \left\{ x(P) + \frac{a}{2} \right\} \chi_{\hat{\tau}}(P).$$

Theorem 14. *For any odd prime p of good reduction,*

$$-\frac{1}{p}S_{\hat{\tau}} = h_p^* + \hat{R}_{a,b}. \quad (22)$$

Proof. By Proposition 6, we first obtain:

$$\begin{aligned}
S_{\hat{\tau}} &= \eta\left(\frac{b}{p}\right) + \sum_{\substack{P \in E_1(\mathbb{F}_p) \\ P \neq T_1, \infty}} \left\{x(P) + \frac{a}{2}\right\} \chi_{\hat{\tau}}(P) \\
&= \eta\left(\frac{b}{p}\right) + \sum_{x=1}^{p-1} \left\{x + \frac{a}{2}\right\} \left(\frac{x}{p}\right) \left(1 + \left(\frac{x^3 + ax^2 + bx}{p}\right)\right) \\
&= \eta\left(\frac{b}{p}\right) + \sum_{x=1}^{p-1} \left\{x + \frac{a}{2}\right\} \left(\frac{x}{p}\right) + \sum_{x=1}^{p-1} \left\{x + \frac{a}{2}\right\} \left(\frac{x^2 + ax + b}{p}\right) \\
&= \sum_{x=1}^{p-1} \left\{x + \frac{a}{2}\right\} \left(\frac{x}{p}\right) + \sum_{x=0}^{p-1} \left\{x + \frac{a}{2}\right\} \left(\frac{x^2 + ax + b}{p}\right).
\end{aligned}$$

The first sum evaluates as follows:

$$\begin{aligned}
\sum_{x=1}^{p-1} \left\{x + \frac{a}{2}\right\} \left(\frac{x}{p}\right) &= \sum_{x=1}^{p-1-\eta} (\{x\} + \{\eta\}) \left(\frac{x}{p}\right) + \sum_{x=p-\eta}^{p-1} (\{x\} + \{\eta\} - p) \left(\frac{x}{p}\right) \\
&= \sum_{x=1}^{p-1} x \left(\frac{x}{p}\right) + \eta \sum_{x=1}^{p-1} \left(\frac{x}{p}\right) - p \sum_{x=p-\eta}^{p-1} \left(\frac{x}{p}\right) = -ph_p^* - p \sum_{x=1}^{\eta} \left(\frac{-x}{p}\right).
\end{aligned}$$

As for the second sum, we substitute $u = x + \eta$ and apply Lemma 7 to get:

$$\sum_{x=0}^{p-1} \left\{x + \frac{a}{2}\right\} \left(\frac{x^2 + ax + b}{p}\right) = \sum_{u=0}^{p-1} \{u\} \left(\frac{u^2 - \frac{r}{4}}{p}\right) = -p\delta_{-r/4} = -p\delta_{-r}.$$

Combining the above calculations yields the result. $\square$

We conclude with a brief comment on the “globalization” of the error term for $S_{\hat{\tau}}$, analogously to Theorem 9 for S_{τ} . Take E_1 , E_2 , and τ as in (7), with $a, b \in \mathbb{Z}$. For simplicity, suppose $p > a > 0$, so that $a = \{a\}$. If a is even, we have the explicit bound $|\hat{R}_{a,b}| \leq \frac{a}{2} + 1$ for the error term. If a is odd, the behavior of the error is different. For $p > 2a$, we have $\eta = \frac{p-1}{2} + \frac{a+1}{2}$ exactly, and so the error term has roughly $\frac{p}{2}$ terms. Further, from [1, Ch. 5, §4], we have

$$\sum_{x=1}^{\frac{p-1}{2}} \left(\frac{-x}{p}\right) = \begin{cases} 0 & p \equiv 1, 5 \pmod{8} \\ -3h_p & p \equiv 3 \pmod{8} \\ -h_p & p \equiv 7 \pmod{8}. \end{cases}$$

Thus, the term $\hat{R}_{a,b}$ may dwarf h_p ! Rather, if we define

$$\hat{\rho}_{a,b} := \delta_{-r} + \sum_{x=1}^{\frac{a+1}{2}} \left(\frac{-\frac{p-1}{2} - x}{p}\right),$$

then $|\hat{\rho}_{a,b}| \leq \frac{a+3}{2}$, and (22) can be rewritten:

$$-\frac{1}{p}S_{\hat{\tau}} = \hat{\rho}_{a,b} + \begin{cases} 0 & p \not\equiv 3 \pmod{8} \\ -2h_p & p \equiv 3 \pmod{8}. \end{cases}$$

This gives a better description of the behavior of the sum $S_{\hat{\tau}}$ when a is odd.

Acknowledgements

We are grateful to Kirti Joshi for first bringing our attention to the interesting divisibility properties of weighted character sums on elliptic curves, and for many fruitful conversations during our research. We give our thanks to Joe Silverman and Matt Papanikolas for helpful suggestions. We wish to recognize the contribution of **Sage** [9], which was invaluable for computational experimentation related to this project. Finally, we appreciate the many helpful comments and suggestions of the referee during the revision of this article.

References

- [1] Z.I. Borevich and I.R. Shafarevich. *Number Theory*, volume 20 of *Pure and Applied Mathematics*. Academic Press, New York, 1966. Translated by Newcomb Greenleaf.
- [2] Harold Davenport. *Multiplicative number theory*, volume 74 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 2000. Revised and with a preface by Hugh L. Montgomery.
- [3] Max Deuring. Die Typen der Multiplikatorenringe elliptischer Funktionenkörper. *Abh. Math. Sem. Hansischen Univ.*, 14:197–272, 1941.
- [4] A. Joux and F. Morain. Sur les sommes de caractères liées aux courbes elliptiques à multiplication complexe. *J. Number Theory*, 55(1):108–128, 1995.
- [5] Hugh L. Montgomery and Robert C. Vaughan. *Multiplicative number theory. I. Classical theory*, volume 97 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2007.
- [6] R. Padma and S. Venkataraman. Elliptic curves with complex multiplication and a character sum. *J. Number Theory*, 61(2):274–282, 1996.
- [7] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1986.
- [8] Joseph H. Silverman. *Advanced topics in the arithmetic of elliptic curves*, volume 151 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994.

- [9] W.A. Stein et al. *Sage Mathematics Software (Version 4.3.3)*. The Sage Development Team, 2009. <http://www.sagemath.org>.
- [10] Kenneth S. Williams. Evaluation of character sums connected with elliptic curves. *Proc. Amer. Math. Soc.*, 73(3):291–299, 1979.